

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Insecta Mundi

Center for Systematic Entomology, Gainesville,
Florida

2016

Classification, natural history, and evolution of the subfamily Peloniinae Opitz (Coleoptera: Cleroidea: Cleridae). Part VI. New taxonomic placement for *Pelonium sexpunctatum* Kirsch

Weston Opitz

Florida State Collection of Arthropods, opitz@kwu.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>

 Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

Opitz, Weston, "Classification, natural history, and evolution of the subfamily Peloniinae Opitz (Coleoptera: Cleroidea: Cleridae). Part VI. New taxonomic placement for *Pelonium sexpunctatum* Kirsch" (2016). *Insecta Mundi*. 1001.

<http://digitalcommons.unl.edu/insectamundi/1001>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0496

Classification, natural history, and evolution of the subfamily Peloniinae
Opitz (Coleoptera: Cleroidea: Cleridae). Part VI. New taxonomic
placement for *Pelonium sexpunctatum* Kirsch

Weston Opitz
Florida State Collection of Arthropods
Florida Department of Agriculture and Consumer Services
P.O. Box 147100
Gainesville, Florida 32614-7100, USA

Date of Issue: July 29, 2016

Weston Opitz

Classification, natural history, and evolution of the subfamily Peloniinae Opitz
(Coleoptera: Cleroidea: Cleridae). Part VI. New taxonomic placement
for *Pelonium sexpunctatum* Kirsch
Insecta Mundi 0496: 1-5

ZooBank Registered: LSID: urn:lsid:zoobank.org:pub:FF884A29-3CE3-4BD0-B124-9F9E7F8A36D8

Published in 2016 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://www.centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology. Manuscript preparation guidelines are available at the CSE website.

Chief Editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Assistant Editor: David Plotkin

Head Layout Editor: Eugenio H. Nearn

Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas

Review Editors: Listed on the Insecta Mundi webpage

Manuscript Preparation Guidelines and Submission Requirements available on the Insecta Mundi web-page at: <http://centerforsystematicentomology.org/insectamundi/>

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and Agrifood Canada, Ottawa, ON, Canada
The Natural History Museum, London, Great Britain
Muzeum i Instytut Zoologii PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Michael C. Thomas

Classification, natural history, and evolution of the subfamily Peloniinae Opitz (Coleoptera: Cleroidea: Cleridae). Part VI. New taxonomic placement for *Pelonium sexpunctatum* Kirsch

Weston Opitz

Florida State Collection of Arthropods

Florida Department of Agriculture and Consumer Services

P.O. Box 147100

Gainesville, Florida 32614-7100, USA

opitz@kwu.edu

Abstract. *Pelonium sexpunctatum* Kirsch (Coleoptera: Cleroidea: Cleridae) is transferred to *Diutius* Opitz where it is closely allied to *Diutius galleruoides* Spinola.

Key words. Taxonomy, species, nomenclatural changes.

Resumen. *Pelonium sexpunctatum* Kirsch (Coleoptera: Cleroidea: Cleridae) es transferido al género *Diutius* Opitz donde es cecado a *Diutius galleruoides* Spinola.

Palabras clave. Taxonomía, especies, cambios de nomenclatura.

Introduction

During a revision of *Pelonium* Spinola (Opitz, in preparation), I received the primary type of *Pelonium sexpunctatum* Kirsch. A “?” placed behind the genus name *Pelonium* by Kirsch (1874: 398) suggests that he had some doubts about the appropriate generic placement of this species. Indeed, the presence of a tibial spur formula of 0-2-2 clearly places *Pelonium sexpunctatum* Kirsch in the genus *Diutius* Opitz. I follow the higher classification proposal for the Cleridae as outlined by Opitz (2010: 56).

Materials and methods

This study is based mostly on two specimens. The primary type of *Pelonium sexpunctatum* Kirsch was received from Olaf Jaeger, Museum of Zoology, Senckenberg Natural History Collections, Dresden, Coleoptera Collection, Königsbruecker Landstrasse 159, D-01109, Dresden Germany (SMTD). The second specimen, a non-type, comes from Antoine Mantilleri, Museum d’Histoire Naturelle, Entomologie, 45 bis, Rue de Buffon, Paris (Ve), France (MNHN). Methods involving measurements and morphological terminology follow those described in Opitz (2010). Abbreviations used in this work are defined as follows: EW/FW= eye width (frontal view)/frons width (frontal view); PW/PL= pronotal width (from apex of one tubercle to apex of the other) /pronotal length (from midline anterior margin to posterior margin); EL/EW= elytral length (from humeral angle to apex)/elytral width (greatest dorsal width of one elytron). Measurements were made at 250X. Microscopic observations were made with a M5 Wild stereoscopic microscope with camera lucida attachment (Leica, Wetzlar, Germany). Habitus photographs were taken with a Leica Z 16 APO microscope equipped with JVC KY-F75U-CCD camera and controlled by Syncroscopy Auto Montage software (Cambridge, United Kingdom).

Taxonomy

Diutius sexpunctatus (Kirsch)

Figures 1, 2, 4, 5.

Pelonium?sexpunctatum Kirsch, 1874: 398. Lectotype. Female. PERU, Pozuzu, M. Kirsch (SMTD). Corporaal 1950: 290 (*Corinthiscus*). Kirsch did not specify as to the number of specimens that were

involved in his description. Therefore, I invoke Recommendation 73F of the ICZN (1999) and designate a lectotype for this nominal species.

Diagnosis. Specimens of this species resemble superficially those of *Diutius galleruoides* (Spinola, 1844: 372), but in *Diutius sexpunctatus* (Kirsch) specimens the elytral disc shows 6 maculae (Fig. 5) not 8 (Fig. 6) as is the case in its congener.

Redescription. *Size:* Length 10.0 mm; width 3.2 mm. *Form:* Oblong, short, rectangular (Fig. 5). *Color:* Mouthparts testaceous, except mandibles brown; antenna mostly testaceous, posterior margin of capitulum brown; thorax, legs, and abdomen testaceous; elytra mostly testaceous, with 6 brown maculae. *Head:* Funicular antennomeres progressively shorter, capitulum longer than combined length of funicular antennomeres, capitular antennomeres very long (Fig. 1); eyes coarsely faceted and bulgy, frons slightly narrower than width of eye (EW/FW 22/20). *Thorax:* Pronotum (Fig. 2) slightly transverse (PW/PL 72/70), side margin with shallow tubercle, disc finely sparsely punctate; elytral without asetiferous punctation (EL/EW 240/60). *Abdomen:* Pygidium scutiform.

Variation. *Size:* Length 9.0-10.0 mm; width 3.0- 3.2 mm. Other than body size, the available specimens are quite homogeneous.

Distribution. In addition to the lectotype, I examined one specimen from: **Peru:** Tarapoto, Mai a Aut, 1886, M de Mathan. Specimens are deposited in MNHN and SMTD.

Discussion

Heretofore, the genus *Diutius* was comprised of two species: *Diutius galleruoides* (Fig. 6) and *D. pallidus* (Fig. 7, 8) (Opitz, 2014: 426). The additional species, *Diutius sexpunctatus* (Fig. 5), may be conveniently included in a key (key to *Akonesis*, *Antennosus*, *Crusbatus*, *Diutius*, and their species) provided by Opitz (2014: 412). To incorporate *Diutius sexpunctatus* into this key, the key should be modified as follows:

- | | | |
|-------|---|---------------------------------------|
| 3(1). | Tibial spur formula 1-2-2 (genus <i>Akonesis</i>) | 4 |
| — | Tibial spur formula 0-2-2 (genus <i>Diutius</i>) | 5 |
| 4(3). | Elytral disc with brown vitta; humeral margin not infuscated (Bolivia to Brazil) | <i>Akonesis vittatis</i> (Spinola) |
| — | Elytral disc without brown vitta, humeral margin with black streak, disc with small black markings (Brazil) | <i>Akonesis pictipennis</i> (Gorham) |
| 5(3). | Elytral disc unicolorous, light brown, without brown maculae (Bolivia) | <i>Diutius pallidus</i> Opitz |
| — | Elytral disc bicolorous, disc with light brown maculations | 6 |
| 6(5). | Elytral disc with 6 maculations (Peru) (Fig. 5) | <i>Diutius sexpunctatus</i> (Kirsch) |
| — | Elytral disc with 8 maculations (Guyana, French Guiana, Ecuador, Brazil) (Fig. 6) | <i>Diutius galleruoides</i> (Spinola) |

The possession of elytral disc maculae is interpreted herein as a synapotypic characteristic, which unites *Diutius sexpunctatus* (Kirsch) and *D. galleruoides* (Spinola) into a sister-group relationship. Together, these two species and *Diutius pallidus* Opitz form a monophyletic group based on the synapotypic characteristic, tibial spur formula 0-2-2.


Acknowledgments

I thank Olaf Jaeger and Antoine Mantilleri for entrusting me with specimens in their charge. John M. Leavengood, Jr., Lionel Stange, and Michael Thomas kindly reviewed the manuscript and my wife, Galena, provided the habitus illustrations. I also thank Paul Skelley for various courtesies afforded me as a Research Associate of The Florida State Collection of Arthropods, Gainesville, Florida. As always, I am grateful to Jean-Michel Maes for the Spanish abstract.


Literature Cited

- Corporaal, J. B. 1950.** Cleridae. In: W.D. Hinks (ed). *Coleopterorum catalogus supplementa*, Pars 23 (editio secunda). W. Junk; The Hague. Netherlands. 373 p.
- (ICZN) International Code of Zoological Nomenclature. 1999.** 4th Edition. Adopted by the International Union of Biological Sciences. The International Trust for Zoological Nomenclature, London, UK., 306 p.
- Kirsch, T. F. W. 1874.** Beiträge zur Kenntniss der Peruanischen Käferfauna auf Dr. Abendroth's Sammlungen basirt. *Berliner Entomologische Zeitschrift* 17: 338–418.
- Opitz, W. 2010.** Classification, natural history, phylogeny, and subfamily composition of the Cleridae and generic content of the subfamilies (Coleoptera: Cleroidea). *Entomologica Basiliensia et Collectionis Frey* 32: 31–128.
- Opitz, W. 2014.** Taxonomy of the New World checkered beetle genera *Akonesis* Opitz, new genus, *Antennosus* Opitz, new genus, *Crusbatus* Opitz, new genus, and *Diutius* Opitz, new genus (Coleoptera: Cleridae: Peloniinae). *The Coleopterists Bulletin*. 68(3): 411–428.
- Spinola, M. 1844.** Essai monographique sur les Clérîtes: insectes Coléoptères. Tome 1, Imprimerie des frères Ponthenier, Gênes, Italy.

Received July 6, 2016; Accepted July 21, 2016
Review editor David Plotkin.



Figures 1-4. Structures and map of *Diutius* species. 1) Antenna of *D. seipunctatus*. 2) Pronotum of *D. seipunctatus*. 3) Pronotum of *D. gallerucoides*. 4) Geographic distribution of *Diutius* species as noted.



Figures 5-8. Habitus and male genitalia. **5)** Habitus of *D. sexpunctatus*. **6)** Habitus of *D. galleruoides*. **7)** Habitus of *D. pallidus*. **8)** Male genitalia of *D. pallidus*.

